

	<i>Testing for Vector-borne Infections</i>	
	<b>Policy Number:</b> POL-PP-313 AHS – G2158	<b>Original Creation Date:</b> 7/1/2025
	<b>Version Number:</b> 002	<b>Version Effective Date:</b> 7/1/2025
	<b>Policy Status:</b> Active	<b>Next Review Date:</b> 7/1/2026

**NOTICE**

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Kansas City (Blue KC).

Blue KC reserves the right to review and revise these policies when necessary. When there is an update, we will publish the most current policy to: <https://providers.bluekc.com/ContactUs/PaymentPolicies>.

PROVIDER/ENTITY IMPACTED					
<input checked="" type="checkbox"/> <b>PROFESSIONAL</b>	<input checked="" type="checkbox"/> <b>FACILITY</b>	<input type="checkbox"/> <b>DME</b>	<input type="checkbox"/> <b>AMBULATORY SURGERY</b>	<input checked="" type="checkbox"/> <b>LAB</b>	<input type="checkbox"/> <b>OTHER</b>

LINES OF BUSINESS IMPACTED						
<input checked="" type="checkbox"/> <b>COMMERCIAL</b>	<input checked="" type="checkbox"/> <b>BLUE MEDICARE ADVANTAGE</b>	<input checked="" type="checkbox"/> <b>ACA QHP<sup>1</sup></b>	<input checked="" type="checkbox"/> <b>SMALL GROUP ACA</b>	<input checked="" type="checkbox"/> <b>JAA<sup>2</sup></b>	<input checked="" type="checkbox"/> <b>FEP<sup>3</sup></b>	<input type="checkbox"/> <b>DENTAL</b>

<sup>1</sup> ACA QHP: Affordable Care Act Qualified Health Plan for Individual/Family    <sup>2</sup> JAA: Joint Administrative Account    <sup>3</sup> FEP: Federal Employee Program

**Disclaimer**

Blue KC has developed Provider Payment Policies to provide guidance on payment methodologies as they pertain to submitted claims. These policies are written following industry standard recommendations from sources such as:

- Current Procedural Terminology
- Centers for Medicare and Medicaid
- American Medical Association
- National Correct Coding Initiative
- Other professional organizations and societies

Coverage of any service is determined by date of service, a member's eligibility and benefit limits for the service or services rendered, all terms of the Provider Service Agreement, and other standards of coding rules and guidelines.

Final payment is subject to the application of claims adjudication and edits common to the industry.

For confirmation of which services may be eligible for coverage and description of when medical services are considered medically necessary, not medically necessary, or investigational, please contact:

- Blue KC Provider Hotline for Commercial lines of Business 816-395-3929
- Affordable Care Act Provider Hotline 866-859-3822
- Blue Medicare Advantage Provider Hotline 866-508-7140

In the event of a conflict between any policies, the Member's coverage document will govern.

## Description/Application

Arthropod vectors, including mosquitoes, ticks, fleas, and mites, that feed on vertebrate hosts can spread bacteria, protozoa, and viruses during feeding to their susceptible host, resulting in a variety of infections and diseases. Arboviruses (arthropod-borne viruses) include Zika virus, West Nile virus (WNV), chikungunya virus, dengue virus (DENV), yellow fever virus (YFV), and Colorado tick fever virus (CTF) to name a few. Malaria and babesiosis are both conditions caused by arthropod-borne protozoan parasites, Plasmodium and Babesia, respectively. Conditions caused by arthropod-borne bacteria include rickettsial diseases, ehrlichiosis, anaplasmosis, and Lyme disease, as well as other Borrelia-associated disorders. Isolation, identification, and characterization of these various infections depend on the causative agent. Identification methods may include culture testing, microscopy, and staining techniques; moreover, molecular testing, such as nucleic acid amplification testing (NAAT), and serologic testing, including immunofluorescence antibody assays and enzyme-linked immunosorbent assays (ELISA), can be used for laboratory diagnosis.

For Lyme disease and testing for *Borrelia burgdorferi*, please see AHS-G2143 Lyme Disease Testing.

## Policy

Application of coverage criteria is dependent upon an individual's benefit coverage at the time of the request.

For individuals suspected of having babesiosis (see Note 1), the use of a Giemsa- or Wright-stain of a blood smear, nucleic acid amplification testing (NAAT), or IgG or IgM indirect immunofluorescence antibody (IFA) assay for Babesia (initial testing and confirmatory testing should occur a minimum of 2 weeks apart) **may be reimbursed**.

For individuals suspected of having relapsing fever caused by *Borrelia* spp, the following testing **may be reimbursed**:

For individuals suspected of having a relapsing fever caused by a *Borrelia* spp., the following testing **may be reimbursed**:

- For individuals suspected of having hard tick relapsing fever (HTRF) (see Note 2): serologic assays to detect *Borrelia* antibodies or NAAT testing to detect *Borrelia miyamotoi*.
- For individuals suspected of having louse-borne relapsing fever (LBRF) (see Note 3): peripheral blood smear microscopy or NAAT testing to detect *Borrelia recurrentis*.
- For individuals suspected of having a soft tick relapsing fever (STRF)/tickborne relapsing fever (TBRF) (see Note 4): dark-field microscopy of a peripheral blood smear, microscopy of a Wright- or Giemsa-stained blood smear, NAAT testing to detect *Borrelia* spp., or serologic assays to detect *Borrelia* antibodies.

For individuals suspected of having a relapsing fever caused by a *Borrelia* spp., culture testing for *Borrelia* **may not be reimbursed**.

For individuals suspected of having chikungunya (see Note 5), the use of viral culture for diagnosis, NAAT for the presence of chikungunya in a blood sample, or IFA assay for IgM antibodies during both the acute and convalescent phases **may be reimbursed**.

For individuals suspected of having Colorado tick fever (CTF) (see Note 6), the use of NAAT testing or IFA for CTF-specific IgM antibodies **may be reimbursed**.

For the detection of dengue virus (DENV), the use of NAAT, IgM antibody capture ELISA (MAC-ELISA), or NS1 ELISA, as well as a confirmatory plaque reduction neutralization test for DENV, **may be reimbursed** in the following individuals:

- For individuals suspected of having a DENV infection (see Note 7).
- For individuals who are symptomatic for Zika virus infection (see Note 8).

For individuals suspected of having DENV (see Note 7), the use of IgG ELISA or hemagglutination testing **may not be reimbursed**.

For individuals suspected of having ehrlichiosis and/or anaplasmosis (see Note 8), the use of NAAT of whole blood, IFA assay for IgG antibodies, or microscopy for morulae detection **may be reimbursed**.

For individuals suspected of having ehrlichiosis and/or anaplasmosis (see Note 8), the use of an IFA assay for IgM antibodies or standard blood culture **may not be reimbursed**.

For individuals suspected of having malaria (see Note 10), the use of a rapid immunochromatographic diagnostic test or smear microscopy to diagnose malaria, determine the species of *Plasmodium*, identify the parasitic life-cycle stage, and/or quantify the parasitemia (can be repeated up to three times within three days if initial microscopy is negative in suspected cases of malaria) **may be reimbursed**.

To confirm the species of *Plasmodium* in an individual diagnosed with malaria, NAAT testing **may be reimbursed**.

The use of IFA for *Plasmodium* antibodies **may not be reimbursed**.

For individuals suspected of having a rickettsial disease (see Note 11), the use of an IFA assay for IgG antibodies (initial testing and confirmatory testing should occur a minimum of two weeks apart) **may be reimbursed**.

For individuals suspected of having a rickettsial disease (see Note 11), the use of standard blood culture, NAAT, or IFA assay for IgM antibodies **may not be reimbursed**.

For individuals suspected of having West Nile virus (WNV) disease (see Note 12), the use of IFA for WNV-specific IgG or IgM antibodies in either serum or CSF and a confirmatory plaque reduction neutralization test for WNV **may be reimbursed**.

To confirm a WNV infection in individuals who are immunocompromised, nucleic acid detection of WNV **may be reimbursed**.

For immunocompetent individuals suspected of having WNV disease (see Note 12), the use of NAAT for WNV **may not be reimbursed**.

For individuals suspected of having a yellow fever virus (YFV) infection (see Note 13), the use of NAAT for YFV or serologic assays to detect virus-specific IgM and IgG antibodies, as well as a confirmatory plaque reduction neutralization test for YFV, **may be reimbursed**.

For the detection of Zika virus, the use of NAAT **may be reimbursed** in the following individuals:

- Up to 12 weeks after the onset of symptoms for symptomatic (see Note 8) pregnant individuals who, during pregnancy, have **either** lived in or traveled to areas with current or past Zika transmission **or** who have had sex with someone who either lives in or has recently traveled to areas with current or past Zika virus transmission (see Note 14).
- For symptomatic non-pregnant individuals living in or with recent travel to an area with an active CDC Zika Travel Health Notice or an area with current or past Zika virus transmission (see Note 14) when symptoms presented within the last seven days.

Zika virus NAAT and Zika virus IgM testing, as well as a confirmatory plaque reduction neutralization test for Zika, **may be reimbursed** in **any** of the following situations:

- Up to 12 weeks after the onset of symptoms for symptomatic (see Note 8) pregnant individuals who, during pregnancy, have **either** lived in or traveled to areas with an active CDC Zika Travel Health Notice **or** who have had sex with someone who either lives in or has recently traveled to areas with an active CDC Zika Travel Health Notice (see Note 14).
- For pregnant individuals who have a fetus with prenatal ultrasound findings consistent with congenital Zika virus infection (see Note 15).
- For infants born from individuals who, during pregnancy, tested positive for Zika virus.
- For infants born with signs and symptoms of congenital Zika syndrome (see Note 15) and who have a birthing parent who had a possible Zika virus exposure during pregnancy.
- For symptomatic non-pregnant individuals living in or with recent travel to an area with an active CDC Zika Travel Health Notice or an area with current or past Zika virus transmission (see Note 14) when symptoms presented more than seven days prior to testing.

For non-pregnant individuals who have not traveled outside of the United States and its territories and who are symptomatic for Zika virus infection (see Note 8), NAAT and/or IgM testing for Zika detection **may not be reimbursed**.

For asymptomatic individuals, testing for babesiosis, chikungunya virus, CTF, DENV, ehrlichiosis and/or anaplasmosis, malaria, rickettsial disease, TBRF, WNV, YFV, or Zika virus during a general exam without abnormal findings **may not be reimbursed**.

#### NOTES:

**Note 1:** Typical signs and symptoms of babesiosis can include hemolytic anemia, splenomegaly, hepatomegaly, jaundice, and nonspecific flu-like symptoms such as fever, chills, body aches, weakness, and fatigue.

**Note 2:** Typical signs and symptoms of HTRF (caused by *Borrelia miyamotoi*) can include chills or shakes, fatigue, nausea or vomiting, headache, and muscle and joint aches.

**Note 3:** Typical signs and symptoms of LBRF (caused by *Borrelia recurrentis*) can include fever, headache, chills or shakes, muscle and joint aches, and nausea. Though the clinical symptoms of LBRF are similar to STRF, LBRF is usually associated with fewer relapses.

**Note 4:** Typical signs and symptoms of STRF/TBRF (caused by *Borrelia hermsii*, *B. turicatae*, and other *Borrelia* bacteria) can include fever, headache, muscle aches, chills, dizziness, joint pain, nausea and vomiting, appetite loss, and rarely, facial paralysis eye pain or redness, or vision changes.

**Note 5:** Typical signs and symptoms of chikungunya include high fever (>102°F or 39°C), joint pains (usually multiple joints, bilateral, and symmetric), headache, myalgia, arthritis, conjunctivitis, nausea, vomiting, and maculopapular rash.

**Note 6:** Typical signs and symptoms of CTF can include fever, chills, headache, myalgia, malaise, sore throat, vomiting, abdominal pain, and maculopapular or petechial rash.

**Note 7:** Typical signs and symptoms of dengue include fever, headache, retro-orbital eye pain, myalgia, arthralgia, macular or maculopapular rash, petechiae, ecchymosis, purpura, epistaxis, gingival bleeding, hematuria, leukopenia, thrombocytopenia, hyponatremia, elevated AST and ALT, and nausea and/or vomiting.

**Note 8:** Typical signs and symptoms of Zika virus infection can include fever, rash, headache, joint pain, conjunctivitis (red eyes), and muscle pain.

**Note 9:** Typical signs and symptoms of ehrlichiosis and/or anaplasmosis usually begin 5-14 days after an infected tick bite, and they include fever, headache, malaise, myalgia, and shaking chills. Ehrlichiosis can also present with gastrointestinal issues, including nausea, vomiting, and diarrhea.

**Note 10:** Typical signs and symptoms of malaria can include fever, influenza-like symptoms (e.g., chills, headache, body aches), anemia, jaundice, seizures, mental confusion, kidney failure, and acute respiratory distress syndrome.

**Note 11:** Typical signs and symptoms of rickettsial diseases (including Rocky Mountain spotted fever, *Rickettsia parkeri* rickettsiosis, *Rickettsia* species 364D rickettsiosis, *Rickettsia* spp. (mild spotted fever), and *R. akari* (rickettsialpox)) usually begin 3 – 12 days after initial bite and can include fever, headache, chills, malaise, myalgia, nausea, vomiting, abdominal pain, photophobia, anorexia, and skin rash. *Rickettsia* species 364d rickettsiosis can also present with an ulcerative lesion with regional lymphadenopathy.

**Note 12:** Typical signs and symptoms of WNV include headache, myalgia, arthralgia, gastrointestinal symptoms, and maculopapular rash. Less than 1% of infected individuals develop neuroinvasive WNV with symptoms of meningitis, encephalitis, or acute flaccid paralysis.

**Note 13:** Typical signs and symptoms of yellow fever include symptoms of the toxic form of the disease (jaundice, hemorrhagic symptoms, and multisystem organ failure), as well as nonspecific influenza symptoms (fever, chills, headache, backache, myalgia, prostration, nausea, and vomiting in initial illness).

**Note 14:** The CDC provides information on the geographic risk classifications of Zika (<https://www.cdc.gov/zika/geo/index.html>), as well as providing travel health notices for pathogens of concern (<https://wwwnc.cdc.gov/travel/notices>).

**Note 15:** Typical signs and symptoms of congenital Zika syndrome can include microcephaly, problems with brain development, feeding problems (e.g., difficulty swallowing), hearing loss, seizures, vision problems, decreased joint movement (i.e., contractures), and stiff muscles (making it difficult to move).

Coding	
CPT	Code Description
86280	Hemagglutination inhibition test (HAI)
86382	Neutralization test, viral
86619	Antibody; Borrelia (relapsing fever)
86666	Antibody; Ehrlichia
86750	Antibody; Plasmodium (malaria)
86753	Antibody: protozoa, not elsewhere specified
86757	Antibody; Rickettsia
86788	Antibody; West Nile virus, IgM
86789	Antibody; West Nile virus
86790	Antibody: virus, not elsewhere specified
86794	Antibody; Zika virus, IgM
87040	Culture, bacterial; blood, aerobic, with isolation and presumptive identification of isolates (includes anaerobic culture, if appropriate)
87164	Dark field examination, any source, (e.g., penile, vaginal, oral, skin) includes specimen collection
87166	Dark field examination, any source, (e.g., penile, vaginal, oral, skin) without specimen collection
87207	Smear, primary source with interpretation; special stain for inclusion bodies or parasites (eg, malaria, coccidia, microsporidia, trypanosomes, herpes viruses)
87449	Infectious agent antigen detection by immunoassay technique, (eg, enzyme immunoassay [EIA], enzyme-linked immunosorbent assay [ELISA], fluorescence immunoassay [FIA], immunochemiluminometric assay [IMCA]) qualitative or semiquantitative; not otherwise specified, each organism
87468	Infectious agent detection by nucleic acid (DNA or RNA); Anaplasma phagocytophilum, amplified probe technique
87469	Infectious agent detection by nucleic acid (DNA or RNA); Babesia microti, amplified probe technique
87478	Infectious agent detection by nucleic acid (DNA or RNA); Borrelia miyamotoi, amplified probe technique
87484	Infectious agent detection by nucleic acid (DNA or RNA); Ehrlichia chaffeensis, amplified probe technique
87662	Infectious agent detection by nucleic acid (DNA or RNA); Zika virus, amplified probe technique

87798	Infectious agent detection by nucleic acid (DNA or RNA), not otherwise specified; amplified probe technique, each organism
87899	Infectious agent antigen detection by immunoassay with direct optical (ie, visual) observation; not otherwise specified
0043U	Tick-borne relapsing fever Borrelia group, antibody detection to 4 recombinant protein groups, by immunoblot, IgM Proprietary test: Tick-Borne Relapsing Fever Borrelia (TBRF) ImmunoBlots IgM Test Lab/Manufacturer: IGeneX Inc
0044U	Tick-borne relapsing fever Borrelia group, antibody detection to 4 recombinant protein groups, by immunoblot, IgG Proprietary test: Tick-Borne Relapsing Fever Borrelia (TBRF) ImmunoBlots IgG Test Lab/Manufacturer: IGeneX Inc

**References and Resources**

Avalon Medical Policy AHS – G2158 – Testing for Vector-borne Infections

**Related Documents**

Avalon Medical Policy AHS- G2143 -Lyme Disease Testing

Avalon Medical Policy AHS -M2097 - Identification of Microorganisms Using Nucleic Acid Probes

**Revision History**

<b>Version</b>	<b>Date</b>	<b>Summary of Revisions</b>
001	07/01/2025	Initial version
002	10/01/2025	Avalon 4th Quarter updates